



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup> :</b>  <b>A61K 7/48</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 00/09082</b>  <b>(43) International Publication Date:</b> 24 February 2000 (24.02.00)
<b>(21) International Application Number:</b> PCT/EP99/05028  <b>(22) International Filing Date:</b> 14 July 1999 (14.07.99)  <b>(30) Priority Data:</b> 14 August 1998 (14.08.98)      GB 9817817.1  <b>(71) Applicant (for AE AU BB CA CY GB GD GH GM IE IL KE LC LK LS MN MW NZ SD SG SL SZ TT UG ZA ZW only):</b> UNILEVER PLC [GB/GB]; Unilever House, Blackfriars, London EC4P 4BQ (GB).  <b>(71) Applicant (for all designated States except AE AU BB CA CY GB GD GH GM IE IL IN KE LC LK LS MN MW NZ SD SG SL SZ TT UG ZA ZW):</b> UNILEVER NV. [NL/NL]; Weena 455, NL-3013 AL Rotterdam (NL).  <b>(71) Applicant (for IN only):</b> HINDUSTAN LEVER LIMITED [IN/IN]; Hindustan Lever House, 165/166 Backbay Reclamation, Mumbai 400 020, Maharashtra (IN).  <b>(72) Inventor:</b> METHMANUS-SPALTRO, Suree; Unilever Thai Holdings Limited, Lever House, 411 Srinakarin Road, Suanlaung, Bangkok 10250 (TH).		<b>(74) Agent:</b> ELLIOTT, Peter, William; Unilever plc, Patent Dept., Colworth House, Sharnbrook, Bedford MK44 1LQ (GB).  <b>(81) Designated States:</b> AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> COSMETIC COMPOSITION  <b>(57) Abstract</b>  An anhydrous foaming cleansing composition for topical application to the human skin, comprising an anionic surfactant, glycerine, polyethylene glycol, and a water insoluble benefit agent.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

## COSMETIC COMPOSITION

This invention relates to cosmetic applications for topical applications to  
5 the human skin. In particular, it relates to anhydrous cosmetic  
compositions, with the capacity for self warming on application.

Anhydrous cosmetic compositions which provide a warming benefit on  
application are as such known. For example, EP-A-27730 (Johnson  
10 Company) describes a cosmetic composition, which may for example be a  
hair treatment or hand treatment composition, which generates heat on  
contact with water. In this application, heat generation is alleged to be  
provided by the presence of polyethylene oxides, polypropylene oxides,  
and derivatives thereof, and a 2 to 9 carbon atom alkaline glycol.  
15 However, we have found that the formulation of compositions based on or  
containing significant amounts of propylene glycol needs to be very  
carefully controlled, since a number of propylene glycol based  
compositions suffer from product separation on storage.

20 In terms of further teachings in the field, EP-A-586929 (Kao) describes a  
two pack system which generates heat on the addition of water, and  
produces a physiologically compatible salt that generates heat on mixture  
with water. Also contributing to the heat felt by the user on hydration of  
the product is the heat of hydration of polyethylene glycol.

25

Further, GB-A-1357000 (British American Tobacco Company) describes a  
topical cosmetic composition comprising an anhydrous polyol and an

absorbent particulate filler material. Suitable polyols are said to include propylene glycol, glycerol, 1,3 butylene glycol, and polyethylene glycols having an average molecular weight of from 2 to 300 and from 1000 to 6000.

5

We have found that a particularly suitable cosmetic composition can be prepared for topical application, which composition is self heating, has good mildness and foaming properties, and which has further benefits not appreciated in the prior art.

10

Thus, according to a first aspect of the invention, there is provided an anhydrous foaming cleansing composition for topical application to the human skin, comprising an anionic surfactant, glycerine, polyethylene glycol, and a water insoluble benefit agent.

15

The surfactant contained in the composition may be any combination of surfactants, provided that it comprises an anionic surfactant and it provides the composition with a suitable level of foaming and mildness.

As such, the surfactant component of the composition may comprise in addition to anionic surfactants soaps, cationic, nonionic, zwitterionic and amphoteric surfactants, and mixtures thereof.

20

Suitable soaps include these having carbon chain lengths of C<sub>8</sub>-C<sub>24</sub>, be saturated or unsaturated, and have any appropriate cation, such as sodium, potassium, ammonium or triethylammonium.

25

The composition according to the invention comprises an anionic surfactant, which is preferably chosen from alkyl sulphates, alkyl ether sulphates, alkyl sulphonates, alkyl aryl sulphonates, olefin sulphonates, acyl sarcosinates, acyl taurides, acyl isethionates, nonoalkyl  
5 sulphosuccinates, dialkylsulphosuccinates, N-acylated  $\alpha$ -amino acids, alkyl carboxylates, monoalkyl phosphates and dialkyl phosphates, and mixtures thereof. Specific examples of suitable anionic surfactants include:

alkyl sulphates, such as sodium lauryl sulphate [eg EMPICOL CX  
10 available from Albright and Wilson], and triethanolamine lauryl sulphate [eg EMPICOL TL40/T, available from Albright and Wilson].

alkylether sulphates, such as sodium lauryl ether sulphate [eg EMPICOL  
ESB70, available from Albright and Wilson].

15


alkyl sulphonates, such as sodium alkane (C<sub>13-18</sub>) sulphonate [eg  
HOSTAPUR SAS 30, available from Hoechst].

alkylaryl sulphonates, such as sodium alkyl benzene sulphonate [eg  
20 TEEPOL CM44, available from Shell].

olefin sulphonates, such as sodium olefin sulphonate (C<sub>5-18</sub>) [eg  
HOSTAPUR OS, available from Hoechst].

25 acyl sarcosinates, having the structure: (51)

4




(51)

- 5 where  $\text{R}^3$  is chosen from  $\text{C}_{6-14}$  alkyl, and  
 M is a counterion chosen from alkali metals, ammonium and substituted  
 ammonium such as alkanolammonium.

An example of an acyl sarcosinate having the structure (51), is sodium  
 10 lauryl sarcosinate [eg HAMPSOYL L-95, available from Grace].

acyl taurides, having the structure (52):




(52)

wherein  $\text{R}^4$  is chosen from  $\text{C}_{8-18}$  alkyl

- 20 An example of an acyl tauride having the structure (52) is coconut methyl  
 taurine [eg FENOPEN TC 42, available from International Specialty  
 Products].

acyl isethionates, having the structure (53):



(53)


wherein R<sup>5</sup> is chosen from C<sub>8-18</sub> alkyl.

5

An example of an acyl isethionate having the structure (53) is sodium acyl isethionate [eg JORDAPON C1, available from Jordon).

monoalkyl sulphosuccinates, having the structure (54):

10



(54)

15 where R<sup>6</sup> is chosen from C<sub>10-20</sub> alkyl.

Examples of monoalkyl sulphosuccinates having the structure (54) include:

20 sodium lauryl sulphosuccinate [eg EMPICOL SLL, available from Albright and Wilson].

magnesium alkyl sulphosuccinate [eg ELFANOL 616 Mg, available from Akzo].

25 sodium lauryl ethoxysulphosuccinate [eg EMPICOL SDD, available from Albright and Wilson].

coconut monoethanolamide ethoxysulphosuccinate [eg EMPICOL SGG].

disodium lauryl polyglycolether sulphosuccinate [eg SURTAGENE S30, available from CHEM-Y].

5

polyethyleneglycol sulphosuccinate [eg REWOPOL SBFA 30, available from REWO].

dialkyl sulphosuccinates, having the structure (55):

10



where  $\text{R}^7$  and  $\text{R}^8$  are the same or different, and are chosen from  $\text{C}_6\text{-}_{14}$  alkyl.

15

An example of a dialkyl sulphosuccinate having the structure (55) is sodium dilauryl sulphosuccinate [eg EMCOL 4500, available from Witco].

20

N-acylated  $\alpha$ -amino acids, such as sodium lauroyl glutamate [eg ACYLGLUTAMTATE LS-11, available from Ajinomoto Co Inc].

alkyl ether carboxylates, such as  $\text{C}_{12-14}\text{O}(\text{EO})_4\text{OCH}_2\text{CO}_2\text{Na}$  [eg AKYPO RLM 38, available from Akzo].

25

monoalkyl phosphates and dialkyl phosphates, such as dioctyl phosphate.

Further examples of anionic surfactants (and of the other types of surfactants) are described in "Surface Active Agents and Detergents" (vols. I and II), by Schwartz, Ferry and Bergh.

In certain embodiments, preferred anionic surfactants include alkyl ether sulphates, fatty acid soaps, alkyl sulphates, alkyl sulphonates, isethionic acid derivatives, and mixtures thereof. In certain embodiments, preferred non-soap anionic surfactants may be C<sub>8</sub>-C<sub>22</sub> alkyl unsubstituted isethionates.

The composition of the invention can also comprise a cationic surfactant. Suitable cationic surfactants are those with the structure (57):



where R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> each represents alkyl or aryl groups, and X represents a halogen counterion.

Preferred cationic surfactants in accordance with structure (57) include:

Hexadecyl trimethyl ammonium chloride, such as Arquad 16, available from Akzo.

Dihydrogenated tallow dimethyl ammonium chloride, such as Arquade 2HT, available from Akzo.


Dodecyl benzyl dimethyl ammonium bromide, such as Amoxyl Br1244,  
5 available from Seppic.

Cocoamidopropyl trimethyl ammonium chloride, such as Empigen CSC, available from Albright and Wilson.

10 The composition of the invention can also comprise an amphoteric surfactant. Suitable amphoteric surfactants are derivatives of aliphatic quaternary ammonium, phosphonium and sulphonium compounds, wherein the aliphatic radicals contain from 8 to 18 carbon atoms, and may be straight chain or branched, and further contain an anionic water  
15 solubilising group, such as carboxyl, sulphonate, sulphate, phosphate or phosphonate.

Preferred amphoteric surfactants include:

20 Alkyl betaines, having the structure (58):




(51)

25 where R<sup>1</sup> is C<sub>1-16</sub> alkyl.

An example of an alkyl betaine having the structure (58) is lauryldimethyl betain [eg EMPIGEN BB, available from Albright and Wilson].


5 Alkylamidopropyl betaines, having the structure (59):



10

An example of an alkylamidopropyl betaine having the structure (59) is cocamidopropyl betaine [eg TEGOBETAIN L7, available from Goldschmidt].

15 Alkylamphoglycinates or Alkylamphopropionates having the structure (60):



20


where  $\text{R}^{11}$  is chosen from H,  $\text{CH}_2\text{COO}^-$  and  $(\text{CH}_2)_2\text{COO}^-$ , and  $\text{R}^{111}$  is chosen from  $\text{CH}_2\text{COO}^-$  and  $(\text{CH}_2)_2\text{COO}^-$

25

Suitable examples of compounds (60) are cocoamphoglycinate (available from International Specialty Products), and cocoamphopropionate.

Sultaines, having the structure (61):

5



10 where R<sup>2</sup> is chosen from C<sub>12-16</sub> alkyl alkylamido groups.

An example of a sultaine having the structure (61) is cocamidopropylhydroxysultaine [eg CYCLOTERIC BET-CS, available from Alcolac).

15

The most preferred amphoteric surfactants are lauryl dimethyl betaine and cocamidopropyl betaine.

Such amphoteric surfactants can contribute to the foaming of the skin  
20 cleansing composition, while ameliorating the harshness of the anionic surfactant.

The composition of the invention can also comprise a nonionic surfactant. Suitable nonionic surfactants include polyoxyethylene alkyl esters,  
25 polyoxyethylene alkyl ethers, and alkyl polyglycosides.

A suitable example of a polyoxyethylene alkyl esters is that having the CTFA designation Polysorbate 80 which is a mixture of oleate esters of sorbitol and sorbitol anhydrides, condensed with approximately 20 moles of ethylene oxide. Also suitable is Polysorbate 20 which is a mixture of  
5 laurate esters or sorbitol and sorbitol anhydrides condensed with approximately 20 moles of ethylene oxide.

Polysorbate 80 and Polysorbate 20 are available commercially as TWEEN 80 and TWEEN 20 respectively, from ICI Americas.

10

Also suitable for use in the compositions of the invention is the polyethylene glycol ether of C<sub>9-11</sub> alcohol with an average of 8 ethoxy units, which is available commercially as NONIDET LE-8T or as SYNPERIONIC 91-8T, and the polyethylene glycol ether of C<sub>12-15</sub> alcohol  
15 with an average of 9 ethoxy units which is available commercially as DOBANOL 25-9.

Particularly useful alkyl polyglycosides include the glycosides of glucose or glucose oligomers where the alkyl chain can be C<sub>8-16</sub> and the average  
20 number of glucose units is 1 to 2. A suitable example is ORAMIX NS 10 which is the glucoside of C<sub>10-12</sub> fatty alcohol with an average of about 1.5 glucose units.

Conveniently the surfactant is present in the composition at a level of 5-  
25 50% by weight. Preferably the level of surfactant in the composition is at least 10% by weight; preferably, the level of surfactant in the composition is less than 35% by weight. The anionic component of the surfactant

content of the composition can typically be 40-100% of the total surfactant content of the composition.

In a preferred embodiment, the composition may contain at least 5%,  
5 more preferably at least 8% by weight, even more preferably at least 10% by weight of sodium cocoyl isethionate surfactant. In such compositions, a distinct improvement in sensory properties of the composition was perceived.

10 Conveniently, the blend of surfactants in compositions according to the invention is selected so as to enhance the lather volume and rinsability of the composition.

Preferably, the polyethylene glycols used in compositions according to the  
15 invention is a blend of polyethylene glycols having a variety of molecular weights; preferably, at least 50% of the polyethylene glycol blend used in compositions according to the invention has a molecular weight of less than 600, conveniently in the molecular weight of 200-600. However, higher molecular weight polyethylene glycols can usefully be used in the  
20 composition provided that they are soluble in the formulation.

Conveniently, the polyethylene glycol used in the composition have a molecular weight of less than 5,000; preferably 95% or more of the polyethylene glycols in the composition have a molecular weight of less  
25 than this.

Compositions according to the invention have been found to have particularly beneficial properties, in terms of their self heating properties on exposure to water, but also in relation to their stability. Further, cosmetic compositions according to the invention which comprise the  
5 blend of glycerine and polyethylene glycol components have been found to be particularly suitable for delivery of components which are not soluble in water, and which may otherwise be difficult to deliver from an aqueous based composition. As such, compositions according to the invention have been found to be particularly suitable for delivery of these  
10 components.

Other benefits which may be attained with compositions according to the invention include the provision of soft radiant skin after use, reduced skin oiliness, moisturization and mildness, as well as good rinsability.

15

An essential component of the composition is glycerine. Preferably, glycerine is present at a level of 5% to 25%, more preferably 10% to 20% by weight of the composition.

20 A further essential component of the composition is a blend of polyethylene glycols, which preferably comprise of at least 50% by weight of the polyethylene glycol component of polyethylene glycols having a molecular weight of 600 or less. The polyethylene glycol component of the composition may also comprise other, high molecular weight  
25 polyethylene glycols, which may typically have a molecular weight up to about 6000. Preferably, the polyethylene glycol component of the

composition comprises from 20% to 70% by weight, more preferably from 25% to 50% by weight of the composition.

5 Compositions according to the invention are anhydrous; that is they contain a maximum of 1% by weight of water.

In certain embodiments, a preferred optional component is propylene glycol, though the level of propylene glycol is carefully controlled, since excessive levels of propylene glycol may prejudice the stability of the  
10 composition. Suitable levels of propylene glycol are typically 0.2-15%, more typically 1-13% by weight of the composition.

As referred to above, compositions according to the invention may be particularly suitable for the delivery of water insoluble components. Such  
15 components include but are not limited to benefit agents such as skin soothing agents, refreshing agents, healing agents, cooling agents, toning agents, blood flow promoting agents, anti-inflammatory agents, moisturising agents, anti-ageing agents, and sunscreen agents. A specific example of a suitable sunscreen agent is Parsol MCX.

20

An advantage of compositions according to the invention is that they may facilitate the solubilization and/or delivery of components which are difficult to deliver from aqueous based compositions, and may be considered insoluble in aqueous compositions. Typically, such benefit  
25 agents may be present at levels of up to 20%, more preferably 0.1-10%, even more preferably 0.5-8% by weight.

The product can take any convenient product form, and is preferably a liquid product. Conveniently, the product can be a facial foam or a shower cream. Conveniently the composition is not encapsulated, but is a free flowing liquid.

5

Products according to the invention have been found to provide suitable skin warming on application, which causes the pores in the skin to open more widely, and thus promote deeper cleansing.

- 10 The invention will now be described by way of example only, with reference to the following examples.

**Example 1**

The following composition provided a suitable self warming facial foam composition according to the invention.

5

<u>Component</u>	<u>%</u>	<u>Tradename</u>	<u>Supplier</u>
Propylene glycol	13.0	propylene glycol	Dow Chemicals
Glycerine	14.0	glycerol	GBT
Potassium laurate/ 10 myristate/stearate	22.0	Nonsoul LK-2 MK1/SK-1	NOF Corp
PEG 300/400/600	29.0	Carbowax 300-600	Union Carbide
PEG 60000 Distearate	3.0	Nionion DS-60HN	Nippon Oils & Fats
15 Ethylene glycol monostearate	8.0	Cutina EGMS	Henkel
20 Oleyl alcohol 20 mole ethoxylate	5.5	Rhodasurf ON-870	Rhone Poulent
Stearic acid	1.0	Pristerene 4900	Unichema
25 Sedaplant extract	1.0	Sedaplant Richter	Chemisches Laboratorium Dr Kurt Richter GmbH
30 Minor ingredients (perfume, etc.)	to 100.0		

**Example 2**

The following composition provided a suitable self warming facial foam composition according to the invention.

5

<u>Component</u>	<u>%</u>	<u>Tradename</u>	<u>Supplier</u>
Propylene glycol	13.0	Propylene Glycol	Dow Chemicals
Glycerine	16.0	Glycerol	LBT
10 Potassium laurate	7.0	Nonsoul LK-2	NAF Corp.
Sodium cocoyl isethionate	11.0	Elfan AT 84G	Akzo
15 PEG 300/400/600	29.0	Carbowax 300/400/600	Union Carbide
PEG 6000 Distearate	3.0	Nionion DS-60HN	Nippon Oil & Fats
20 Ethylene glycol monostearate	8.0	Cutina EGMS	Henkel
Oleyl alcohol 25 20 mole ethoxylate	6.0	Rhodasurf ON-870	Rhone Poulenc
Stearic acid	1.1	Pristerine 4900	Unichema
30 Sedaplant extract	1.0	Sedaplant Richter	Chemisches Laboratorium Dr Kurt Richter GmbH
35 Minor ingredients (perfume, etc.)	to 100		

## CLAIMS

1. An anhydrous foaming cleansing composition for topical application to the human skin, comprising an anionic surfactant, glycerine, polyethylene glycol, and a water insoluble benefit agent.  
5
2. A cleansing composition according to claim 1, wherein the anionic surfactant is a C<sub>8-22</sub> alkyl unsubstituted isethionate.
- 10 3. A cleansing composition according to claim 2, comprising at least 5% of sodium cocoyl isethionate.
4. A cleansing composition according to any of the preceding claims, wherein the surfactant comprises 5-50% by weight of the composition.  
15
5. A cleansing composition according to claim 4, wherein the composition comprises at least 10% by weight of surfactant.
6. A cleansing composition according to any of the preceding claims, wherein the anionic surfactant comprises 40-100% by weight of the surfactants in the composition.  
20
7. A cleansing composition according to any of the preceding claims, wherein at least 50% by weight of the polyethylene glycol in the composition has a molecular weight of 200-600.  
25

8. A cleansing composition according to any of the preceding claims, wherein the polyethylene glycol in the composition has a molecular weight of less than 5,000.
- 5 9. A cleansing composition according to any of the preceding claims, wherein the glycerine comprises 5-25% by weight of the composition.
- 10 10. A cleansing composition according to any of the preceding claims, wherein the polyethylene glycol comprises 20-70% by weight of the composition.
11. A cleansing composition according to any of the preceding claims, wherein the composition contains less than 1 % by weight water.
- 15 12. A cleansing composition according to any of the preceding claims, wherein the composition additionally comprises 0.2-15% by weight propylene glycol.
- 20 13. A cleansing composition according to any of the preceding claims, wherein the composition additionally comprises an amphoteric surfactant.
14. A cleansing composition according to claim 13, wherein the amphoteric surfactant comprises at least 5% by weight of the composition.
- 25 15. A cleansing composition according to any of the preceding claims, wherein the benefit agent is selected from as skin soothing agents, refreshing agents, healing agents, cooling agents, toning agents, blood

flow promoting agents, anti-inflammatory agents, moisturising agents, anti-ageing agents, and sunscreen agents.

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 99/05028

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K7/48

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	DE 196 24 870 A (HENKEL KGAA) 2 January 1998 (1998-01-02) ---	
A	EP 0 261 754 A (SCHERER LTD R P) 30 March 1988 (1988-03-30) ---	
A	PATENT ABSTRACTS OF JAPAN vol. 1998, no. 02, 30 January 1998 (1998-01-30) & JP 09 272894 A (S T CHEM CO LTD;GOING:KK), 21 October 1997 (1997-10-21) abstract --- -/--	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

### \* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

29 October 1999

Date of mailing of the international search report

08/11/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Stienon, P

# INTERNATIONAL SEARCH REPORT

In ternational Application No

PCT/EP 99/05028

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	PATENT ABSTRACTS OF JAPAN vol. 017, no. 052 (C-1022), 2 February 1993 (1993-02-02) & JP 04 264186 A (OSAKA AEROSOL IND CORP), 18 September 1992 (1992-09-18) abstract ---	
A	EP 0 027 730 A (JOHNSON CO LTD) 29 April 1981 (1981-04-29) cited in the application ---	
A	EP 0 586 929 A (KAO CORP) 16 March 1994 (1994-03-16) cited in the application ---	
A	GB 1 357 000 A (BRITISH AMERICAN TOBACCO CO) 19 June 1974 (1974-06-19) cited in the application -----	

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 99/05028

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DE 19624870 A	02-01-1998	AU 3257497 A WO 9749374 A EP 0910327 A NO 985963 A	14-01-1998 31-12-1997 28-04-1999 18-12-1998
EP 0261754 A	30-03-1988	GB 2192404 A AT 67086 T DE 3772915 A	13-01-1988 15-09-1991 17-10-1991
JP 09272894 A	21-10-1997	NONE	
JP 04264186 A	18-09-1992	NONE	
EP 0027730 A	29-04-1981	JP 56057711 A JP 56077216 A AU 6346880 A	20-05-1981 25-06-1981 02-07-1981
EP 0586929 A	16-03-1994	DE 4227203 A AT 110955 T DE 69300008 D DE 69300008 T JP 7069835 A US 5538720 A	07-10-1993 15-09-1994 13-10-1994 26-01-1995 14-03-1995 23-07-1996
GB 1357000 A	19-06-1974	DE 2145032 A FR 2106584 A IE 36942 B ZA 7106230 A	06-04-1972 05-05-1972 30-03-1977 28-06-1972